

Total Marks: 80

Time Duration: 3Hr

N.B.: 1) Question no.1 is compulsory.

2) Attempt any three questions from Q.2 to Q.6.

3) Figures to the right indicate full marks.

Maximum

Marks

- Q1. a)** Find the Laplace transform of $\cos 2t \sin t e^{-t}$. [5]
- b)** Find the half-range sine series for $f(x) = x(\pi - x)$ in $(0, \pi)$. [5]
- c)** Show that the function $f(z) = ze^z$ is analytic and find $f'(z)$ in terms of z . [5]
- b)** Prove that $\nabla \left\{ \nabla \cdot \frac{\vec{r}}{r} \right\} = -\frac{2}{r^3} \vec{r}$. [5]
- Q2. a)** Find the inverse Z-transform of $F(z) = \frac{z}{(z-1)(z-2)}$, $|z| > 2$. [6]
- b)** Find the analytic function whose real part is $\frac{\sin 2x}{\cosh 2y + \cos 2x}$. [6]
- c)** Obtain Fourier series for the function $f(x) = \begin{cases} 1 + \frac{2x}{\pi}, & -\pi \leq x \leq 0 \\ 1 - \frac{2x}{\pi}, & 0 \leq x \leq \pi \end{cases}$, [8]
deduce that $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$
- Q3. a)** Find $L^{-1} \left[\frac{1}{s^2(s+a)^2} \right]$ using convolution theorem. [6]
- b)** Show that the set of functions $\cos nx$, $n = 1, 2, 3, \dots$ is orthogonal on $[0, 2\pi]$. [6]
- c)** Using Green's theorem evaluate $\int_C \left(\frac{1}{y} dx + \frac{1}{x} dy \right)$ where C is the boundary of the region defined by $x = 1, x = 4, y = 1$ and $y = \sqrt{x}$. [8]
- Q4. a)** Find Laplace transform of $f(t) = k \frac{t}{T}$ for $0 < t < T$ and $f(t) = f(t + T)$. [6]
- b)** Show that $\vec{f} = (x^2 + xy^2) \vec{i} + (y^2 + x^2y) \vec{j}$ is irrotational and find its scalar potential. [6]
- c)** Find half-range cosine series for $f(x) = x$, $0 < x < 2$. Using Parseval's identity deduce that [8]
i) $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} - \frac{1}{5^4} + \dots$
ii) $\frac{\pi^4}{90} = \frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \dots$
- Q5. a)** Use divergence theorem to show that $\iint_S \nabla r^2 \cdot \vec{ds} = 6v$ where S is any closed surface enclosing a volume V . [6]
- b)** Find the Z-transform of $f(k) = k\alpha^k$, $k \geq 0$. [6]
- c)** i) Find $L^{-1} \left[\frac{(s+2)^2}{(s^2+4s+8)^2} \right]$ [8]
ii) Find $L^{-1} [2 \tanh^{-1} s]$
- Q6. a)** Solve using Laplace transform [6]
 $(D^2 - 3D + 2)y = 4e^{2t}$, with $y(0) = -3, y'(0) = 5$.
- b)** Find the bilinear transformation which maps the points 1, -i, 2 on z -plane onto 0, 2, -i respectively of w -plane. [6]
- c)** Express the function $f(x) = \begin{cases} \sin x, & 0 < x \leq \pi \\ 0, & x < 0, x > \pi \end{cases}$ as Fourier integral and deduce [8]
that $\int_0^\infty \frac{\cos\left(\frac{w\pi}{2}\right)}{1-w^2} dw = \frac{\pi}{2}$.
